

ON THE SHAPE OF PULSE SPECTRA IN GAMMA-RAY BURSTS

FELIX RYDE AND ROLAND SVENSSON

Stockholm Observatory, S-133 36 Saltsjöbaden, Sweden

ABSTRACT

The discovery (Liang & Kargatis 1996), that the peak energy of time-resolved spectra of gamma-ray burst (GRB) pulses decays exponentially with fluence, is analytically shown to imply that the time-integrated photon number spectrum of a pulse should have a unique shape, given by an underlying E^{-1} behavior. We also show that the asymptotic low energy normalization of the time-integrated spectrum is equal to the exponential decay constant. We study analytically how this general behavior is modified in more realistic situations and show that diversity is then introduced in the properties of time-integrated GRB pulse spectra. We argue that further diversity will occur in time-integrated multi-pulse (complex) GRB spectra. The total energy received per cm^2 is approximately the decay constant times the maximum peak energy of the pulse. Our analytical results connect the properties of the time-integrated pulse spectrum with those of the time-resolved spectra, and can thus be used when studying observed GRB pulse spectra. We illustrate with the bright burst GRB 910807 and comment on GRB 910525 and GRB 921207.

Subject headings: gamma rays: bursts

1. INTRODUCTION

The time-integrated and time-resolved spectra of gamma-ray bursts (GRB) differ. The general appearance of time-integrated GRB spectra is that of a smoothly broken power law. Time-integrated photon number spectra from missions prior to the *Compton Gamma-Ray Observatory* (CGRO) with its Burst and Transient Source Experiment (BATSE) were often fitted with an “optically thin thermal bremsstrahlung” (OTTB) spectrum, $\mathcal{N}_E(E) \propto E^{-1} e^{-E/E_0}$, combined with a power law at larger energies. Band et al. (1993) showed that nearly all time-integrated BATSE spectra were well fitted by the “Band” function consisting of two power laws, $\mathcal{N}_E(E) \propto E^\alpha$ at low energies and $\mathcal{N}_E(E) \propto E^\beta$ at high energies, smoothly joined by an exponential roll-over characterized by a constant E_0 . The Band function index, α , shows a broad peak between -1.25 and -0.25 (Band et al. 1993), instead of the earlier claimed universal value of -1, β clusters fairly narrowly around -2.1 (Preece et al. 1997), while E_0 has values between 100 keV and 1 MeV, but mainly clustering around 100-200 keV (Band et al. 1993).

Time-resolved spectra, even within single pulses, show strong time evolution, and are, in general, harder than time-integrated spectra (e.g., Pendleton et al. 1994a, Ford et al. 1995, Liang & Kargatis 1996, hereafter LK96, Crider et al. 1997a, 1997b). Most importantly, LK96 found that the peak energy, E_{pk} , of the time-resolved $E^2 \mathcal{N}_E(E)$ spectra of a single pulse in a number of bright and smooth GRBs, decays exponentially as a function of photon fluence, i.e.,

$$E_{\text{pk}}(t) = E_{\text{pk,max}} e^{-\Phi(t)/\Phi_0}. \quad (1)$$

where $E_{\text{pk,max}}$ is the maximum value of E_{pk} within a pulse, $\Phi(t)$ is the photon fluence integrated from the time of $E_{\text{pk,max}}$, and Φ_0 is the exponential decay constant. Furthermore, Crider et al. (1997a) found the instantaneous Band function index, α_i , in general to be hard, > -1 , reaching values as large as +1.6.

The aim of this paper is to show the consequences of the time evolution of $E_{\text{pk}}(t)$, for the time-integrated GRB spectrum of a single pulse, as well as for a number of pulses, and

to determine the meaning of Φ_0 . The following notation is used: $N(t)$ (photons $\text{cm}^{-2} \text{s}^{-1}$) is the photon number flux of the burst at time t , and $N_E(E, t)$ (photons $\text{cm}^{-2} \text{s}^{-1} \text{keV}^{-1}$) is the instantaneous spectrum of the burst at photon energy E (keV) at time t . Here, $N_E(E, t)dE$ is the photon number flux at time t within an energy interval dE around E . Furthermore, $\mathcal{N}_E(E) = \int_{t_1}^{t_2} N_E(E, t) dt$ (photons $\text{cm}^{-2} \text{keV}^{-1}$) is the time-integrated spectrum of the burst and E_{pk} is the peak photon energy of $E^2 \mathcal{N}_E(E, t)$ spectra. The total photon number fluence is $\Phi(t) = \int_0^t N(t') dt'$ (cm^{-2}). Note that $d\Phi(t)/dt = N(t)$.

In § 2, we show analytically that the exponential E_{pk} -decay implies that the time-integrated spectrum of a single pulse should have a typical asymptotic low energy behavior of $\mathcal{N}_E(E) \approx \Phi_0 E^{-1}$, i.e., the same low energy power law slope as the OTTB spectrum (the “universal” $\alpha = -1$), and with the asymptotic low energy normalization given by Φ_0 . Modifications to this behavior below $E_{\text{pk,max}}$ are shown to be approximately given by an incomplete gamma function. In § 3, we discuss the implications for the overall spectra of GRBs and in § 4, we illustrate our analytical findings by analyzing spectral data from the bright burst, GRB 910807. In particular, we show that the decay constant, Φ_0 , of the spectral evolution is consistent with the asymptotic normalization of the time-integrated spectrum. Finally, in § 5, we discuss the effects of a time-varying spectral shape and illustrate using GRB 950525 and GRB 921207

2. THE TIME-INTEGRATED SPECTRUM OF A SINGLE PULSE

We present a scenario in which the shape of the time-integrated spectrum is a direct consequence, and therefore a probe of, the specific time-evolution of the instantaneous spectra. We assume the latter to be peaked in the EN_E spectrum, i.e., have a peak energy E_p around which the number of emitted photons dominate, which is equivalent to that the power law indices $\alpha > -1$ and $\beta < -1$. This is indeed the case in most studied pulses (see, e.g., Crider et al. 1997a), except at the very

end of the pulse decay, when slopes with $\alpha < -1$ sometimes are observed. This does not necessarily contradict the following discussion, as it can be ascribed to the possible presence of a soft component (Preece et al. 1996) and the contributions of previous (possibly untriggered or unresolved) pulses. It should also be noted that the determination of α becomes more difficult the closer the peak energy is to the low energy boundary of the observed spectral range. In any case, such a soft contribution should be negligible compared to the total fluence for the discussion below to be useful.

Integrating the instantaneous spectra over some time interval (t_1 to t_2) of the pulse, the time-integrated spectrum becomes

$$\begin{aligned}\mathcal{N}_E(E) &= \int_{t_1}^{t_2} N_E(E, t) dt = \\ &= \int_{E_{\text{pk}, \text{max}}}^{E_{\text{pk}, \text{min}}} N_E(E, t(E_{\text{pk}})) \frac{dt}{dE_{\text{pk}}} dE_{\text{pk}}, \quad (2)\end{aligned}$$

where $E_{\text{pk}, \text{max}} = E_{\text{pk}}(t = t_1)$ and $E_{\text{pk}, \text{min}} = E_{\text{pk}}(t = t_2)$. If Equation (1) is valid during the time interval being considered, we have

$$\frac{dE_{\text{pk}}(t)}{dt} = -E_{\text{pk}}(t) \frac{N(t)}{\Phi_0} = -E_{\text{pk}}(t) \frac{\hat{N}(E_{\text{pk}}(t))}{\Phi_0}, \quad (3)$$

where we introduce a new function, $\hat{N}(E_{\text{pk}}(t)) = N(t)$. We note that the energy dependence of $N_E(E, t(E_{\text{pk}}))$ normally is on the ratio E/E_{pk} , but that its spectral shape could be time dependent, adding additional dependence on $t(E_{\text{pk}})$ or E_{pk} . We therefore define a function $\hat{N}_E(E/E_{\text{pk}}, E_{\text{pk}})$, where $N_E(E, t) = \hat{N}_E(E/E_{\text{pk}}, E_{\text{pk}}) = \hat{N}(E_{\text{pk}}) f_E(E/E_{\text{pk}}, E_{\text{pk}})$, and where the function $f_E(E/E_{\text{pk}}, E_{\text{pk}})$ describing the instantaneous spectral shape is normalized to unity, $\int f_E(E/E_{\text{pk}}, E_{\text{pk}}) dE = 1$. Then the time-integrated spectrum, Equation (2), becomes

$$\mathcal{N}_E(E) = \Phi_0 \int_{E_{\text{pk}, \text{min}}}^{E_{\text{pk}, \text{max}}} f_E(E/E_{\text{pk}}, E_{\text{pk}}) \frac{dE_{\text{pk}}}{E_{\text{pk}}}. \quad (4)$$

Introducing $f_x(x, E_{\text{pk}}) \equiv f_E(E/E_{\text{pk}}, E_{\text{pk}}) dE/dx = E_{\text{pk}} f_E(E/E_{\text{pk}}, E_{\text{pk}})$, where $x \equiv E/E_{\text{pk}}$, and $f_x(x, E_{\text{pk}})$ is normalized to unity, $\int f_x(x, E_{\text{pk}}) dx = 1$, we find that

$$\mathcal{N}_E(E) = \frac{\Phi_0}{E} \cdot I(E), \quad (5)$$

where the correction factor

$$I(E) \equiv \int_{x_{\text{min}}(E)}^{x_{\text{max}}(E)} f_x(x, E/x) dx, \quad (6)$$

and where $x_{\text{min}}(E) = E/E_{\text{pk}, \text{max}}$ and $x_{\text{max}}(E) = E/E_{\text{pk}, \text{min}}$. We show below that the E -dependence of $I(E)$ is rather weak between $E_{\text{pk}, \text{min}}$ and $E_{\text{pk}, \text{max}}$ for the cases we consider. The first factor in (5), describing the over-all spectral shape, shows that the time-integrated spectrum of a single pulse, which has an exponentially decaying peak energy, is a power law with a time-integrated spectral index of $\alpha_{\text{ti}} \sim -1$ between $E_{\text{pk}, \text{min}}$ and $E_{\text{pk}, \text{max}}$ normalized by the decay constant Φ_0 . This result points out how the instantaneous and the time-integrated spectra are related. It shows that the exponential decay behavior of the instantaneous spectra leaves a signature on the time-integrated spectrum: the decay constant, Φ_0 ,

is its asymptotic low energy normalization and the “universal” time-integrated $\alpha_{\text{ti}} = -1$ is obtained. The only assumption made for the derivation is the empirical relation (1) shown so far to be valid for numerous burst pulses (LK96, Crider et al. 1997a,c), but whose exact validity must be explored.

Equation (1) means that a linear increase in fluence by equal steps of Φ_0 photons cm^{-2} corresponds to a decrease of $\ln E_p$ in equal logarithmic steps. The instantaneous spectra are, roughly, dominated by the about Φ_0 photons cm^{-2} in a logarithmic interval around $E = E_{\text{pk}}$, and as a result $EN_E(E)$ is a constant $= \Phi_0$. As $EN_E(E)$ also is the time-integrated specific flux spectrum, it is clear that the photon spectrum, $\mathcal{N}_E(E)$ of a single pulse has a slope of -1 . The OTTB-like spectrum is thus a result of the *exponential* decay of the peak energy versus photon fluence.

Let us now study the behavior of $I(E)$ when we can neglect the time-dependent changes of the instantaneous spectral shape, i.e., when the E/x dependence of $f_x(x, E/x)$ in Equation (6) can be neglected. It has been shown for several bursts that the time-resolved spectra are well fitted by the Band function and therefore it is of interest to use the Band function in (6). However, before discussing these results, we study two conceptually simpler approximations, which are valuable in illustrating the resulting behavior of the time-integrated spectrum. Firstly, in the case of integrating over all energies, i.e., $x_{\text{min}} = 0$ and $x_{\text{max}} = \infty$, $I(E) = 1$ (due to the normalization of $f_x(x)$) and $\mathcal{N}_E(E) = \Phi_0/E$ is an exact solution of (5). The shape of the instantaneous spectra does not affect the normalization and shape of the time-integrated spectrum as long as the instantaneous $EN_E(E, t)$ spectra are peaked with the Band indices $\alpha_i > -1$ and $\beta_i < -1$. Then the dominant contribution to the integral comes from $x \sim 1$ and $I(E)$ is convergent. Secondly, if we consider finite integration limits in (6) and replace the instantaneous spectra with a Dirac delta function at $E = E_{\text{pk}}$, i.e., $x = 1$ and $f_x(x) = \delta(x - 1)$, we arrive at a similar result: $I(E) = 1$ for $E_{\text{pk}, \text{min}} \leq E \leq E_{\text{pk}, \text{max}}$ and 0 otherwise.

Instantaneous spectra broader than a delta function will introduce deviations from the E^{-1} power law found in the previous case. We now consider the Band function in (6). Changing variables to $y \equiv E/E_0$ and thus $y = x(2 + \alpha_i)$ the Band function becomes

$$f_y(y) = \begin{cases} [A]^{-1} y^{\alpha_i} e^{-y} & \text{if } y < (\alpha_i - \beta_i); \\ [A']^{-1} y^{\beta_i} & \text{if } y \geq (\alpha_i - \beta_i). \end{cases}$$

For f_y to be a continuous function, we have

$$\frac{1}{A'} = \frac{1}{A} (\alpha_i - \beta_i)^{\alpha_i - \beta_i} e^{-(\alpha_i - \beta_i)} \equiv \frac{1}{A} C(\alpha_i - \beta_i), \quad (7)$$

and the normalization, $\int f_y dy = 1$, gives

$$\begin{aligned}A &= \Gamma(\alpha_i + 1) [P(\alpha_i + 1, \alpha_i - \beta_i) - \\ &\quad - \frac{e^{-(\alpha_i - \beta_i)}}{\Gamma(\alpha_i + 1)(\beta_i + 1)} (\alpha_i - \beta_i)^{\alpha_i + 1}]. \quad (8)\end{aligned}$$

where $P(\alpha_i + 1, y)$ is the incomplete gamma function (see, e.g., Press et al. 1992). We have three cases:

(i) $y_{\text{max}} > (\alpha_i - \beta_i)$ and $y_{\text{min}} < (\alpha_i - \beta_i)$ gives

$$\begin{aligned}I(E) &= \frac{\Gamma(\alpha_i + 1)}{A} [P(\alpha_i + 1, \alpha_i - \beta_i) - P(\alpha_i + 1, y_{\text{min}}) \\ &\quad + \frac{C(\alpha_i - \beta_i)}{\Gamma(\alpha_i + 1)(\beta_i + 1)} (y_{\text{max}}^{\beta_i + 1} - (\alpha_i - \beta_i)^{\beta_i + 1})]. \quad (9)\end{aligned}$$




Fig. 1.— The correction factor, $I(E)$, given by Eq. (6) plotted as a function of $E/E_{\text{pk,max}}$ for different values of the instantaneous α_i . The curves are for $f_x(x)$ being a Band function, but the principle behavior below $E_{\text{pk,max}}$ is fairly well described by Equation (10) alone, *i.e.*, for $f_x(x)$ being an exponentially cut-off power law. The solid curves describe a situation with no low energy cut-off, $E_{\text{pk,min}} = 0$. Then Eq. (10) becomes $I(E) = [\Gamma(\alpha_i + 1)/A][1 - P(\alpha_i + 1, y_{\text{min}})]$. The dotted curves include such a cut-off, assuming $E_{\text{pk,max}}/E_{\text{pk,min}} = 20$. This corresponds to using the full Eq. (10).

(ii) $y_{\text{max}} < (\alpha_i - \beta_i)$ gives

$$I(E) = \frac{\Gamma(\alpha_i + 1)}{A} [P(\alpha_i + 1, y_{\text{max}}) - P(\alpha_i + 1, y_{\text{min}})]. \quad (10)$$

This is the spectrum below $(\alpha_i - \beta_i)E_0$, and it is independent of β_i .

(iii) $y_{\text{min}} > (\alpha_i - \beta_i)$ gives

$$I(E) = \frac{\Gamma(\alpha_i + 1)}{A} \cdot \frac{C(\alpha_i - \beta_i)}{\Gamma(\alpha_i + 1)(\beta_i + 1)} (y_{\text{max}}^{\beta_i + 1} - y_{\text{min}}^{\beta_i + 1}). \quad (11)$$

At $y_{\text{max}} = (\alpha_i - \beta_i)$ cases (i) and (ii) are identical, and at $y_{\text{min}} = (\alpha_i - \beta_i)$ cases (i) and (iii) are identical.

Equation (5), together with (9)-(11), relates the shape of the time-integrated spectrum to the five parameters of the instantaneous spectra and their time evolution, α_i , β_i , Φ_0 , $E_{0,\text{max}}$, and $E_{0,\text{min}}$.

The general behavior of the result (9)-(11) is captured well in an approximation where the instantaneous spectra are described as exponentially cut-off power laws, $f_y(y) = [A]^{-1} y^{\alpha_i} e^{-y}$, where $A = \Gamma(\alpha_i + 1)$. This corresponds to a Band function without a high energy power law, *i.e.*, to the case with $\beta_i = -\infty$. The result of the integration using this approximation has the advantage of resulting in the very simple analytical expression, Equation (10), valid at all energies. The effects on the time-integrated spectrum due solely to a finite high energy cut-off energy, $E_{\text{pk,max}}$, is easily seen in (10). By letting $E_{\text{pk,min}} = 0$ and thus $y_{\text{max}} = \infty$, the first term on the right-hand-side of Equation (10) equals unity. Similarly, a non-zero $E_{\text{pk,min}}$ will give rise to deviations from the E^{-1} -law. Asymptotically, at $y_{\text{max}} \ll 1$, *i.e.*, at $E \ll E_{\text{pk,min}}$, the time-

integrated spectrum approaches

$$\mathcal{N}_E(E) = \frac{\Phi_0}{E_{0,\text{min}} \Gamma(\alpha_i + 2)} \left(\frac{E}{E_{0,\text{min}}} \right)^{\alpha_i}, \quad (12)$$

i.e., the time-integrated spectrum takes the low energy shape of the instantaneous spectrum.

The result of integrating instantaneous spectra described by Band functions, *i.e.*, Equations (9)-(11), is shown in Figure 1. The spectra are in the form of the correction factor, $I(E)$, and are plotted against $E/E_{\text{pk,max}}$ for different values of α_i . The time-integrated specific flux spectra, $EN_E(E)$, would be a constant $= \Phi_0$ in the delta function approximation, *i.e.*, $I(E) = 1$. Here, with a more realistic description of the instantaneous spectra, there are deviations from the constant Φ_0 in a boundary interval below $E_{\text{pk,max}}$. The width of this boundary interval depends on the shape of the instantaneous spectra themselves. With an instantaneous photon index of $\alpha_i = -1$, the integrated spectrum will never reach the asymptotic level of Φ_0 . The solid curves in Figure 1 show the effect on the Φ_0/E -spectrum, due to the existence of a maximal peak energy. The dotted curves also include the effect of a low energy bound to the peak energy.

As seen in Figure 1, the existence of a maximum and minimum peak energy makes the time-integrated photon Band-index, α_{ti} , that one would obtain when fitting the spectrum, take a range of values enclosing the “universal” value of -1. This would correspond to part of the “diversity” of α_{ti} seen by Band et al. (1993).

It is useful to introduce $E_p(t)$ as the peak photon energy of the instantaneous $EN_E(E, t)$ spectra as the existence of such a peak corresponds to the assumptions of the presented scenario (*i.e.*, $\alpha_i > -1$ and $\beta_i < -1$). In the case of the Band function, E_{pk} and E_p are related through $E_{\text{pk}} = E_p(2 + \alpha_i)/(1 + \alpha_i)$. The exponential decay found by LK96 holds for both E_p and E_{pk} as they used a constant α_i in their study.

The total time-integrated specific flux (keV cm^{-2}) of a pulse,




Fig. 2.— Flux spectrum of the first pulse in GRB 910807 (467) and the best fit model (solid curve); cf. Fig. 1. See the text for the fitting parameters. The data were rebinned to 10σ . The dotted line shows the asymptotic $\alpha_{ti} = -1$ power law, with the corresponding fitted normalization, Φ_0 . Note that Φ_0 can be directly read off the ordinate. The dashed curve shows the time-integrated spectrum for the same parameters as the best fit model but with $E_{pk,min} = 0$.

$\mathcal{F} = \int_0^\infty E \mathcal{N}_E(E) dE$, is dominated by the Φ_0 photons cm^{-2} in a logarithmic interval near $E_{p,max}$, and we thus expect $\mathcal{F} \approx \Phi_0 E_{p,max}$. Indeed, performing the analytical integrations over E , using $\mathcal{N}_E(E)$ for the delta function case and the exponentially cut-off power law case, both give $\mathcal{F} = \Phi_0 (E_{p,max} - E_{p,min}) \approx \Phi_0 E_{p,max}$. A similar result is obtained for the Band function case when the time-integrated $\beta_{ti} \ll -2$, but as β_{ti} approaches -2 , the high energy power law makes an increasing and non-negligible contribution. The total energy of a pulse coming from an isotropic GRB at a distance d is approximately given by $4\pi d^2 \Phi_0 E_{p,max}$, making Φ_0 a useful quantity for estimating global energetics.

3. MULTI-PULSE SPECTRA OF GRBS

The derived time-integrated spectrum, Equation (5), is for a single pulse within the GRB. Most GRBs, however, consist of several overlapping or resolved pulses. If the $E_{pk,min}$ -values of all pulses are below the minimum energy of the observed spectrum, then all pulses should have a similar low energy slope according to our derivation, but their $E_{pk,max}$ and Φ_0 may differ. The sum of a number of time-integrated pulse spectra will lead to a total spectrum with two spectral “breaks”, one at the $E_{pk,max}$ of the individual pulse with the smallest $E_{pk,max}$, and the second at the largest $E_{pk,max}$. The time-integrated low energy spectrum of k pulses in a burst is the sum of the individual pulse spectra, $\mathcal{N}_{E,tot}(E) = \sum_k \mathcal{N}_{E,k}(E) = E^{-1} \sum_k \Phi_{0,k} I_k(E)$ which would give $\alpha \sim -1$ below the smallest $E_{pk,max}$. This slope may, however, not be observable due to the curvature of the individual pulse spectra (see Fig. 1) and as the smallest $E_{pk,max}$ may be close to or below the minimum energy of the observed spectrum. Between the smallest and the largest $E_{pk,max}$, the spectrum will be a superposition of individual pulse spectra and the spectral shape will be determined by the distributions of Φ_0 and $E_{pk,max}$ among the pulses. If, for instance, Φ_0 is the same for all pulses and their $E_{pk,max}$ is evenly distributed in logarithmic energy, the

spectrum $E^2 \mathcal{N}_E(E)$ will be constant, *i.e.*, the photon index will be ~ -2 . Above the largest $E_{pk,max}$, the spectrum will show the high energy behavior of individual pulses.

The spectral diversity seen when fitting time-integrated spectra of GRBs (Band et al. 1993) is therefore expected to mainly be due to the different statistical distributions of Φ_0 and $E_{pk,max}$ in different GRBs. The spectra of individual pulses, on the other hand, are expected to have a spectral shape similar to the ones shown in Figure 1. Simple GRBs with few pulses may have bumpy spectra (possibly producing previously studied line features, see, *e.g.*, Pendleton et al. 1994b), while complex GRBs with many pulses will have smooth spectra.

4. GRB 910807

The spectra plotted in Figure 1 resemble real GRB pulse spectra. To illustrate this and the discussion above, we wish to study a bright burst, exhibiting a smooth, long, single pulse, as well separated as possible from other pulses. Generally, the first pulse in a burst of several separated pulses should be least contaminated by soft contributions from previous pulses. There are only a few bursts that satisfy these requirements, *e.g.*, the first pulse in the bright GRB 910807 (trigger numbers 647; 3rd BATSE catalogue, Meegan et al. 1996). Furthermore, this burst has very hard time-resolved spectra and no significant evolving spectral shape. The spectra only evolve in the sense that the peak energy decreases. This motivates the use of our approximation in our deduction of the analytical expressions for the time-integrated spectrum, in which we neglected the explicit E_{pk} -dependence.

In Figure 2, we have plotted the time-integrated spectrum of the Large Area Detector (LAD) data for the first pulse of GRB 910807. It is the specific flux spectrum, *i.e.*, $E \mathcal{N}_E(E)$, and therefore the Φ_0 value can directly be read off the ordinate. The dotted line shows the constant Φ_0 -spectrum which is then modified by the $I(E)$ -factor. The solid curve is the best fit of the analytical result (5), with $I(E)$ found by integrating the Band




Fig. 3.— The time-evolution of the shape of the instantaneous spectra of the first pulse in GRB 910807 (467), described by the Band parameters α_i and β_i . The dotted line indicates the power law index -1 . Note that there is only a weak time dependence.

function, *i.e.*, equations (9)-(11) above in §2. The parameters of this fit can be compared to the behavior of the time-resolved spectra. For instance, their exponential decay, described by the decay constant, Φ_0 can be compared to the asymptotic normalization of the time-integrated spectrum.

The best fit to the time-integrated spectrum (see Fig. 2) gives the following parameters: $\Phi_0 = 46 \pm 18 \text{ cm}^{-2}$, $E_{\text{pk,max}} = 300 \pm 40 \text{ keV}$, $E_{\text{pk,min}} = 110 \pm 25 \text{ keV}$, $\alpha_i = 0.43 \pm 0.26$ and $\beta_i = -4.2 \pm 0.8$ with $\chi^2/\text{d.o.f.} = 113.5/113$. The α_i - and β_i -parameters in the fit of the time-integrated spectrum should be interpreted as effective values, *i.e.*, the values they would have had in the absence of any spectral shape-evolution.

The time-resolved spectra were fitted with the Band function with its four parameters: α_i , β_i , E_{pk} , and a normalization. The fits were good, having an averaged $\chi^2/\text{d.o.f.} = 104/110$. Figure 3 presents the values of α_i and β_i of these fits. Note that there is no great variation in the shape of the spectra, *i.e.*, the α_i - and β_i -parameters remain approximately constant, which makes the use of our approximations in §2 relevant. The weighted averages of these two parameters are $\langle \alpha_i \rangle = 0.42$ and $\langle \beta_i \rangle = -2.8$. We numerically checked that a weak spectral shape-evolution, *e.g.*, from $\alpha_i = 0.6$ to 0.2 gives a time-integrated spectrum almost identical to that for a constant $\alpha_i = 0.4$. In search of Φ_0 , we fit the exponential decay law (1) to the measured E_{pk} vs. Φ , and find $\Phi_0 = 50 \pm 4 \text{ cm}^{-2}$. E_{pk} varied from $280 \pm 20 \text{ keV}$ down to $120 \pm 12 \text{ keV}$. This is merely a factor of two, which results in that the boundary effects become important over the whole spectral range.

Note the expected agreement between the fit of the time-integrated spectrum and the parameters from the fits of the time-resolved spectra. The study above has shown that by merely studying the time-integrated spectrum, information about the instantaneous spectra can be extracted. This is made possible by knowing in what way the time-integrated spectrum is formed from the intrinsic spectra.

5. EFFECTS OF A TIME-VARYING SPECTRAL SHAPE

In the discussion above, we assumed that the explicit dependence of $f_x(x, E_{\text{pk}})$ on E_{pk} could be neglected, *i.e.*, the shape of the instantaneous spectra is time-independent. This allowed us to find illustrative analytical expressions describing the time-integrated spectrum. The correction to the general time-integrated spectral shape, due to such a time dependence, is not easily found analytically. It should be pointed out that α_i has been observed to vary with time (see, *e.g.*, Crider et al. 1997a). It is, however, obvious that the effects on the time-integrated spectra should be small for moderately varying pulse shapes, especially in the case of the instantaneous spectra being hard. A comprehensive study must be made numerically, in which the detailed time-evolution of the spectral shape is modeled. Preliminary results show that the analytical expressions found above, *i.e.*, the “averaged” spectrum, differ from the numerically integrated spectra, in which α_i is allowed to vary from its minimal to its maximal value, by 20 – 30 per cent or less in the most cases. To explore how this affects the fitted parameters, we have studied two bursts in which α_i evolves notably and, in particular, approaches -1 . It is also of interest to study bursts, in which the peak energy varies over a larger range, and thus revealing more of the underlying E^{-1} -behavior, *i.e.*, the asymptotic normalization level.

In the first pulse of GRB 920525 (trigger numbers 1625; 3rd BATSE catalogue, Meegan et al. 1996), α_i varies between ~ -0.3 and ~ -1.0 , E_{pk} decays from $1250 \pm 300 \text{ keV}$ to $150 \pm 70 \text{ keV}$, and the Φ_0 -value¹ found from the time-resolved evolution of E_{pk} is $33 \pm 1 \text{ cm}^{-2}$. For details of the spectral evolution, see, *e.g.*, Figure 5 in Ford et al. (1995), Figure 3 in LK96, and Figure 2 in Crider et al. (1997a). The fact that the instantaneous spectra are soft and evolve requires caution to be taken when the analytical models presented above are used. In minimizing the χ^2 of the fit to the data, two equally good minima are found ($\chi^2/\text{d.o.f.} = 97/112$), both having some dif-

¹LK96 found $\sim 20 \text{ cm}^{-2}$ using LAD5, while we use LAD4, which has a stronger signal. We also use a wider energy range. This illustrates that the Φ_0 results are dependent on the details of the analysis and therefore should be treated with some caution.




Fig. 4.— Same as Fig.2., but for the first pulse in GRB 920525 (1625). The large energy interval over which E_{pk} evolves allows the time-integrated spectrum to reach the asymptotic level of Φ_0




Fig. 5.— Same as Fig.3., but for the first pulse in GRB 921207 (2083). At the end of the studied time-interval, α_i approaches -1 .




Fig. 6.— Same as Fig.2., but for the first pulse in GRB 921207 (2083)

difficulties in describing the actual time-evolution. The first fit has $\Phi_0 = 27 \pm 2 \text{ cm}^{-2}$ but the intrinsic spectra are found to be too hard; $\alpha_i = +0.6 \pm 0.5$, and the E_{pk} -values are too small: $E_{\text{pk}} = 800 - 10 \text{ keV}$. The second fit has reasonable values for $\alpha_i = -0.87 \pm 0.05$ and $\beta_i = -2.15 \pm 0.13$, while the rest of the parameters, although consistent with the time-resolved fits, are poorly constrained: $\Phi_0 = 230 \pm 220 \text{ cm}^{-2}$, $E_{\text{pk,max}} = 930 \pm 330 \text{ keV}$ and $E_{\text{pk,min}} = 280 \pm 140 \text{ keV}$. In judging these results, it should be noted that the spectra do have significant spectral evolution, which is not accounted for in the fitting. Furthermore, the instantaneous spectra are quite soft, which makes the constraining of the parameters difficult. Figure 4 shows the time-integrated spectrum of the first pulse in GRB 920525. The large energy range over which the peak energy decays and the low value of $E_{\text{pk,min}}$ allows the spectrum to reach the asymptotic level of Φ_0 .

Another example, in which α_i evolves to soft values is the first pulse in GRB 921207 (trigger numbers 2083; 3rd BATSE catalogue, Meegan et al. 1996; for details, see Figure 5 in Ford et al. 1995, and Figure 1 in Crider et al. 1997c), with α_i varying from ~ 0.0 to ~ -0.7 (as shown in Fig. 5), E_{pk} decaying from $470 \pm 25 \text{ keV}$ to $105 \pm 5 \text{ keV}$ and with a Φ_0 -value of $64 \pm 2 \text{ cm}^{-2}$ found from fitting the time-resolved exponential decay of E_{pk} . As shown in Figure 5 the α_i parameter is constant in the beginning of the evolution but approaches $\alpha_i = -1$ at the end, which could cause problems for the constant- α_i -approach. The time-integrated fit (see Fig. 6), however, correspond quite well to the time-resolved results. The fit to the time-integrated spectrum gives $\alpha_i = 0.0 \pm 0.2$, $\beta_i = -3.4 \pm 0.2$, $E_{\text{pk,max}} = 505 \pm 15 \text{ keV}$ and $E_{\text{pk,min}} = 68 \pm 12 \text{ keV}$, and an asymptotic normalization of $\Phi_0 = 70 \pm 10 \text{ cm}^{-2}$. The relatively good agreement between the time-integrated and the time-resolved fits could suggest that the spectral shape actually is constant in time, or at least that this is a good approximation, and that the late softening is caused by erroneous fits due to the low energy power law disappearing beyond the observed energy range. Note that Crider et al. (1997c) suggest a break in the exponential decay.

6. DISCUSSION

We have presented a scenario in which the exponential decay behavior of the peak energy vs. fluence leads to a time-integrated spectrum with an underlying E^{-1} -behavior. This spectral behavior is not necessarily directly visible in a real spectrum, as it is modified by various effects, such as the existence of upper and lower cut-off energies, $E_{\text{pk,max}}$ and $E_{\text{pk,min}}$. We were able to treat these effects analytically. Other effects, such as a time-varying spectral shape, require numerical treatment and is left for future work.

We derived a five parameter expression for the time-integrated spectrum. These parameters describe the properties of the instantaneous spectra and their time evolution. Fitting a time-integrated spectrum with this expression thus gives information about the time-resolved spectra. We have illustrated our scenario by analyzing spectra from three GRB pulses. In future work we will consider a larger sample of pulses.

Studies of GRB spectra, in particular, those of individual pulses are expected eventually to lead to an understanding of the physical processes responsible for the GRB emission. We emphasize that it is the instantaneous spectra and their time evolution that reflect these processes, rather than the time-integrated spectrum. Unfortunately, most theoretical spectral models assume that it is the time-integrated spectrum that reflects the underlying physical emission mechanism.

We are grateful to Jerry Bonnell for his generous support during our work with the BATSE data and the Wingspan program, and for his hospitality during FR's visit to Goddard Space Flight Center. Furthermore, we are grateful to David Band, Andrei Beloborodov, Claes-Ingvar Björnsson, Gunnlaugur Björnsson, Anthony Crider, Juri Poutanen and Boris Stern for various help. We acknowledge support from the Swedish Natural Science Research Council (NFR) and from a Nordic Project grant at Nordita. FR is indebted to the Royal Swedish Academy of Sciences for travel support.

REFERENCES

- Band, D. L., et al. 1993, *ApJ*, 413, 281
- Crider, A., et al. 1997a, *ApJ*, 479, L39
- Crider, A., Liang, E. P., & Preece, R. D. 1997b, in *Proc. Gamma-Ray Bursts: 4th Huntsville Symposium* (New York: AIP), ed. Meegan, C. A. et al., p.359
- Crider, A., Liang, E. P., & Preece, R. D. 1997c, in *Proc. Gamma-Ray Bursts: 4th Huntsville Symposium* (New York: AIP), ed. Meegan, C. A. et al., p.63
- Ford, L. A., et al. 1995, *ApJ*, 439, 307
- Liang, E. P., & Kargatis, V. E. 1996, *Nature*, 381, 495 (LK96)
- Meegan, C. A., et al. 1996, *ApJS*, 106, 65
- Pendleton, G. N., et al. 1994a, *ApJ*, 431, 416
- Pendleton, G. N., et al. 1994b, in *Proc. 2nd Huntsville Gamma-ray Burst Workshop* (New York: AIP), eds. Fishman, G. J., Brainerd, J. J., & Hurley, K., p.313
- Preece, R. D., Briggs, M. S., Pendleton, G. N., Paciesas, W. S., Matteson, J. L., Band, D. L., Skelton, R. T., & Meegan, C. A. 1996, *ApJ*, 473, 310
- Preece, R. D., Pendleton, G. N., Briggs, M. S., Mallozzi, R. S., Paciesas, W. S., Band, D. L., Matteson, J. L., & Meegan, C. A. 1997, *ApJ*, 496, 849
- Press, W. H., Teukolsky, S. A., Vetterling, W. T., & Flannery, B. P., *Numerical Recipes in Fortran*, 2nd ed., Cambridge: Cambridge Univ. Press, 1992